

学校编码: 10384

学号: 200225053

分类号 _____ 密级 _____

UDC _____

厦 门 大 学
硕 士 学 位 论 文

含铁多相催化剂催化的丙烯环氧化

Epoxidation of Propylene Catalyzed by Iron-Containing
Heterogeneous Catalysts

郭 倩

指导教师姓名: 王 野 教授

专 业 名 称: 物 理 化 学

论文提交日期: 2005 年 7 月

论文答辩时间: 2005 年 月

学位授予日期: 2005 年 月

答辩委员会主席: _____

评 阅 人: _____

2005 年 7 月

A thesis submitted to Xiamen University for M. S. Degree

**Epoxidation of Propylene Catalyzed by Iron-Containing
Heterogeneous Catalysts**

By Qian Guo

Supervisor: Prof. Ye Wang

The State Key Laboratory for Physical Chemistry of Solid
Surface and Department of Chemistry, Xiamen University

July, 2005

厦门大学学位论文原创性声明

兹呈交的学位论文，是本人在导师指导下独立完成的研究成果。本人在论文写作中参考的其他个人或集体的研究成果，均在文中以明确方式标明。本人依法享有和承担由此论文而产生的权利和责任。

声明人（签名）：

年 月 日

摘要	I
Abstract	II
 第一章 绪 论	
1.1 催化与选择氧化反应	1
1.2 丙烯环氧化	2
1.2.1 丙烯氧化热力学分析	3
1.2.2 丙烯气相环氧化	4
1.2.2.1 光催化氧化体系	4
1.2.2.2 Ag 催化体系	4
1.2.2.3. Au 催化体系	6
1.2.3 丙烯液相环氧化	7
1.2.3.1 TS-1 直接作催化剂	7
1.2.3.2 TS-1 负载的贵金属体系	9
1.2.3.3 金属卟啉催化剂	9
1.2.3.4 其它催化体系	10
1.3 分子筛与催化剂设计	10
1.3.1 多孔材料的分类	10
1.3.2 沸石分子筛	12
1.3.3 介孔分子筛	13
1.3.3.1 介孔分子筛的特点	13
1.3.3.2 介孔分子筛活性位的导入	14
1.4 本文的构思、目的和构成	15
参考文献	17

第二章 实验部分

2.1 原料与试剂	21
2.2 MCM-41 催化剂的合成	22
2.2.1 MCM-41 的合成	22
2.2.2 MCM-41 中铁的导入	22
2.2.3 添加碱金属盐 Fe-MCM-41 催化剂的制备	23
2.3 Ferrisilicate 催化剂的合成	24
2.3.1 Ferrisilicate 的合成	24
2.3.2 Ferrisilicate 分子筛的水蒸气处理	24
2.3.3 添加碱金属盐的 ferrisilicate 和 s-ferrisilicate 催化剂的制备	24
2.4 含铁分子筛的表征	25
2.5 催化反应性能测试	26
参考文献	28

第三章 Fe-MCM-41 分子筛的丙烯环氧化反应性能研究

3.1 引言	29
3.2 结果与讨论	30
3.2.1 Fe-MCM-41 催化剂的丙烯环氧化性能	30
3.2.1.1 丙烯气相环氧化的实现	30
3.2.1.2 铁源对催化性能的影响	31
3.2.1.3 含不同阴离子钾盐的影响	32
3.2.1.4 不同合成方法得到的 Fe-MCM-41 催化剂的反应性能	35
3.2.1.5 添加 KCl 后不同 Fe-MCM-41 催化剂的反应性能	36
3.2.1.6 KCl 添加量变化对 Fe-MCM-41 催化剂反应性能的影响	37
3.2.2 Fe-MCM-41 的表征	40
3.2.2.1 XRD 表征	40

3.2.2.2 低温N ₂ 物理吸附	48
3.2.2.3 Raman 光谱表征	53
3.2.2.4 紫外可见漫反射	57
3.2.2.5 H ₂ -TPR结果	63
3.3 催化剂上铁物种结构与反应性能的关联	66
3.4 本章小结	67
参考文献	69

第四章 具有 MFI 结构的 ferrisilicate 微孔分子筛的丙烯环氧化性能研究

4.1 引言	71
4.2 结果与讨论	71
4.2.1 Ferrisilicate 的丙烯环氧化性能	71
4.2.1.1 Ferrisilicate 的反应性能	71
4.2.1.2 水蒸气处理的影响	73
4.2.2 催化剂表征	76
4.2.2.1 XRD 结果	76
4.2.2.2 Raman 结果	81
4.2.2.3 UV-Vis 结果	85
4.2.2.4 H ₂ -TPR结果	90
4.4 本章小结	94
参考文献	95

第五章 结论

5.1 Fe-MCM-41 催化剂	96
5.2 Ferrisilicate 催化剂	97

5.3 钾盐的作用	97
硕士在读期间发表论文目录	98
致谢	101

CONTENTS

Abstract in Chinese	I
Abstract in English	II

Chapter 1 General Introduction

1.1 Catalysis and Selective Oxidation	1
1.2 Epoxidation of Propylene	2
1.2.1 The Analysis in Thermodynamics of Epoxidation of Propylene	3
1.2.2 Gas Epoxidation Systems	4
1.2.2.1 Photoepoxidation Systems	4
1.2.2.2 The Systems with Ag as the Catalysts	4
1.2.2.3. The Systems with Au as the Catalysts	6
1.2.3 Liquid Epoxidation Systems	7
1.2.3.1 The Systems with TS-1 as the Catalysts	7
1.2.3.2 The Systems with Noble Metals Supported by TS-1 as the Catalysts	9
1.2.3.3 The Systems with Metal Porphyrin as the Catalysts	9
1.2.3.4 Other Catalytic Systems	10
1.3 Molecular Sieves and the Catalysts' Design	10
1.3.1 Classification of Molecular Sieves	10
1.3.2 Zeolites	12
1.3.3 Mesoporous Molecular Sieves	13
1.3.3.1 The Characteristic of Mesoporous Molecular Sieves	13
1.3.3.2 The Introduction of active sites to Mesoporous Molecular Sieves	14
1.4 The Objectives and Outline of this Thesis	15
Reference	17

Chapter 2 Experimental

2.1 Materials and Reagents	21
2.2 The Preparation of Fe-MCM-41 Catalysts	22
2.2.1 The Synthesis of MCM-41	22
2.2.2 The Introduction of Iron to MCM-41	22
2.2.3 The Preparation of KCl Modified Fe-MCM-41 Catalysts	23
2.3 The Preparation of ferrisilicate Catalysts	24
2.3.1 The Synthesis of ferrisilicate	24
2.3.2 The Steam Activation Treatment of ferrisilicate	24
2.3.3 The Preparation of KCl added ferrisilicate and s-ferrisilicate	24
2.4 The Characterizations of Iron Containing Molecular Sieves	25
2.5 The Catalytic Reaction	26
References	28

Chapter 3 Epoxidation of Propylene Catalyzed by Fe-MCM-41

3.1 Introduction	29
3.2 Results and Discussion	30
3.2.1 The Catalytic Performances of Fe-MCM-41 for the Propylene Epoxidation	30
3.2.1.1 The Achievements of the Gas Propylene Epoxidation	30
3.2.1.2 The Effects of Iron Resources	31
3.2.1.3 The Effects of Anions	32
3.2.1.4 The Catalytic Performances of Fe-MCM-41 Synthesized by Different Methods	35
3.2.1.5 The Effects of Addition of KCl on Various Fe-MCM-41 ...	36
3.2.1.6 The Effects of the Change in K/Fe on Fe-MCM-41	37
3.2.2 The Characterizations of Fe-MCM-41	40
3.2.2.1 XRD Results	40
3.2.2.2 N ₂ -adsorption Results	48

3.2.2.3 Raman Spectroscopy Results	53
3.2.2.4 UV-Vis DPS Results	57
3.2.2.5 H ₂ -TPR Results	63
3.3 The Relationships of Iron Species Structures	
and the Catalytic Performance	66
3.4 Conclusions	67
References	69

Chapter 4 The Catalytic Performances of ferrisilicate with MFI Structure in Epoxidation of Propylene

4.1 Introduction	71
4.2 Results and Discussion	71
4.2.1 The Catalytic Performances of ferrisilicate	
for the Propylene Epoxidation	71
4.2.1.1 The Performances of ferrisilicate	71
4.2.1.2 The Effects of Steam Activation on ferrisilicate	73
4.2.2 Catalysts Characterizations	76
4.2.2.1 XRD Results	76
4.2.2.2 Raman Results	81
4.2.2.3 UV-Vis DPS Results	85
4.2.2.4 H ₂ -TPR Results	90
4.4 Conclusions	94
References	95

Chapter 5 General Conclusions

5.1 Fe-MCM-41 Catalysts	96
5.2 Ferrisilicate Catalysts	97
5.3 The Effects of Potassium Salts	97

Publication list	98
Acknowledgements	101

厦门大学博士论文摘要库

摘 要

本论文合成了含铁分子筛，考察了分子筛中不同位置的铁物种在丙烯环氧化反应中的催化反应性能差异，并对催化剂结构进行了详细的表征，进而深入探讨了催化剂结构和性能之间的关联。

首先，利用直接水热合成法，模板剂离子交换法和浸渍法合成了铁处于分子筛中不同位置的Fe-MCM-41 催化剂，并向其中添加钾盐进行修饰。研究发现，添加KCl后，不同方法合成的Fe-MCM-41 均能够实现丙烯氧化由烯丙基氧化向环氧化的转变。多种表征手段结果表明，添加KCl会导致骨架外铁物种由扭曲的八面体结构向为 K^+ 离子所稳定的正四面体结构转变，该正四面体结构的铁物种是丙烯环氧化的活性位。对于骨架内四配位的铁物种来说，加入KCl会引起骨架铁物种的脱出，而引起分子筛结构的破坏，但脱出骨架的铁物种由于受到骨架的影响，难以与 K^+ 形成正四面体构型。铁配位结构上的差别导致了其在丙烯气相环氧化中反应性能的不同。对不同阴离子钾盐的考察发现，添加KCl的催化剂具有最好的催化性能。

本论文合成了 ferrisilicate 催化剂，对其进行水蒸汽处理，随后以 KCl 进行修饰，同时以未经水蒸汽处理的催化剂添加 KCl 作为对比，研究其在丙烯环氧化反应中的性能。催化性能结果表明，水蒸汽处理后，添加 KCl 的催化剂在 ferrisilicate 系列催化剂中表现出了最好的催化性能。各种表征结果表明，ferrisilicate 中的铁物种与骨架结合紧密，仅添加 KCl，难以将铁脱出骨架并形成有效的活性位，经水蒸汽处理后，则能成为表面铁物种，在 KCl 的作用下可以形成有利于丙烯进行环氧化反应的正四面体配位结构的新铁物种。

通过对 Fe-MCM-41 和 ferrisilicate 的研究发现，催化剂中添加 KCl 不仅可以使铁物种形成有利于丙烯环氧化反应的正四面体结构，还可以使铁的可还原性能显著降低，从而抑制晶格氧的活性，进而抑制了烯丙基氧化。

关键词：丙烯环氧化；含铁分子筛；KCl 修饰；水蒸汽处理

厦门大学博硕士论文摘要库

Abstract

In this dissertation, KCl modified molecular sieves that contain iron species in- or extra-framework are synthesized to study their different catalytic performances for propylene epoxidation with N_2O . These catalysts have been characterized in details and the structure-reactivity relationships have also been investigated.

Fe-MCM-41 catalysts are synthesized and iron would be introduced into different positions of MCM-41 by direct hydrothermal synthesis (DHT), template-ion exchange (TIE) and impregnation (IMP) methods. It is found that the main reaction route can be shift from the allylic oxidation to the epoxidation over Fe-MCM-41 catalysts with modification by KCl. The characterizations by XRD, N_2 -adsorption, Raman, UV-Vis and H_2 -TPR indicate that after the modification, the coordination structure of ex-frame iron species is probably changed from distorted octahedron to tetrahedron stabilized by potassium ions. The tetrahedrally coordinated iron sites are proposed to be the active sites for the propylene epoxidation. As to the in-frame iron species, they are impelled to migrate into extra-framework positions during the KCl modifying process, which cause the collapse of mesopores. The migrated iron species hardly form tetrahedral conformation with potassium ions because of the impact of framework, which results in the low propylene conversion over KCl-Fe-MCM-41-DHT catalysts. The effect of other potassium salts has been studied. It is found that the catalyst modified by KCl exhibits the best catalytic performances for propylene epoxidation.

A series of different ferrisilicate catalysts are steam-activated and then modified by KCl. Meanwhile, ferrisilicate that are not steam-activated but modified by KCl are used as comparisons to illuminate the effect of different-positioned iron species on propylene epoxidation. Catalysts that are steam-activated and then modified by KCl exhibit superior catalytic performances. The detailed characterizations indicate that in-frame iron species cannot form effective active sites for epoxidation without steam-activated. Steam-activation is required to massively create extra-framework iron species. The coordination structure of resulted extra-framework iron species is changed to tetrahedral coordination that favors epoxidation of propylene. KCl-s-ferrisilicate (Si/Fe=64, K/Fe=2, steam-activated time: 8h) catalyst shows the best catalytic performance.

The studies also found that KCl played the important roles. First, the reducibility of iron species became difficult, indicating the reactivity of lattice oxygen was inhabited and the allylic oxidation was also suppressed. The second is that iron tetrahedral coordination structure was formed and stabilized by KCl, which contribute to epoxidation of propylene. Keywords: Propylene epoxidation, iron-containing molecular sieves, KCl modification, steam-activation

厦门大学博硕士论文摘要库

第一章 绪 论

1.1 催化与选择氧化反应

选择氧化反应在自然界和人类社会占据着重要的地位。催化剂参与了大部分选择氧化反应，如与生命体活动紧密相关的酶催化反应。催化选择氧化是化学品工业生产的重要组成部分。在以催化生产的化学产品中氧化产物占有很高比例。以烷烃类为原料的选择氧化因其资源丰富、价格便宜、产物附加值高等原因，而成为了催化研究的热点研究领域。

Table 1-1 Industrial process for oxidation of hydrocarbons developed during the past 4 decades^[1]

Process	Phase	Catalyst
Naphthalene to phthalic anhydride	Gas	V-K-O
Ethylene to acetaldehyde	Liquid	Pd-Cu-Cl
p-Xylene to terephthalic acid	Liquid	Cu-Mo-Cl
Butene to maleic anhydride	Gas	V-P-O
Methanol to formaldehyde	Gas	Fe-Mo-O
Propylene ammoxidation to acrylonitrile	Gas	Complex oxide
Propylene to acrolein	Gas	Complex oxide
Acrolein to acrylic acid	Gas	V-Mo-O
Ethylene & acetic acid to vinyl acetate	Liquid	Pd-Cu-Cl
o-Xylene to phthalic anhydride	Gas	V-Ti-O
Propylene to propylene oxide	Liquid	Ti-Mo-O
Butanol to methylacrylate	Liquid	MoO ₃
Ethylene to vinyl chloride	Gas	Cu chloride
Butane to maleic anhydride	Gas	V-P-O
Phenol to hydroquinone & catechol	Liquid	TS-1

表 1-1 是近年来应用在烃类选择氧化工业中的反应^[1]，这些反应已经成为整个现代化学工业的一个重要支柱。虽然研究人员在烃类选择氧化方面

Degree papers are in the "[Xiamen University Electronic Theses and Dissertations Database](#)". Full texts are available in the following ways:

1. If your library is a CALIS member libraries, please log on <http://etd.calis.edu.cn/> and submit requests online, or consult the interlibrary loan department in your library.
2. For users of non-CALIS member libraries, please mail to etd@xmu.edu.cn for delivery details.

厦门大学博硕士论文摘要库